SENT BY:2127516849

;11- 2-98 ; 3:45PM ;MORGAN & FINNEGAN\_NY→

202 308 4242;# 3/ 6

#19

			S. DEPARTMENT OF COMMERCE						ATTY, DOCKET NO. 2026-4205US		SIGNAL NO. 08/533,895			
INFORMATION DISCLOSURE CITATION								NO <sup>1</sup>	V 0 3 199	8				
								GROUP 1		) IP 16	2			
(0	Use sevei	rai she	eets ij	nece	ssary	")			APPLICANT(S)					
									Topalian, ct al.					
									FILING DATE September 26, 1995 GROUP ART UNIT 1645					
U.S. PATENT DOCUMENTS														
EXAMINER INITIAL			DOCUMENT NUMBER				ER		DATE	NAME	CLASS	ASS SUBCLASS IF APPROPRIAT		
JW		5	2	6	2	ı	7	7	11/16/93	Brown et al.				
JW		5	3	4	2	7	7	4	08/31/94	Boon et al.				
					_	F	ORE	JGN	PATENT D	OCUMENTS				
	DUCUMENT NUMBER				DATE CO	COUNTRY	CLASS	SUBCLASS	TRANSLA	TIÓN				
													YES	NO
OM		9	4	0	5	3	0	4	3/94	PCT			<del></del>	
		0	6	6	8	3	5	0	08/23/95	EPO				
		3	3	4	1	3	6	7	05/24/84	GERMANY (DE)				+-
		2	1	3	3	5	4	3	08/24/84	GB				<del>  </del>
		9	5	2	3	5	6	7	08/24/95	PCT				
		9	3	2	4	1	8	9	10/13/94 07/22/93	PCT				+-
		9	4	-	4	4	5	9	07/94	PCI			· · · · · · · · · · · · · · · · · · ·	1
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Papers, Etc.)  Chiez, R.M., et al "Specificity And Promiscuity Among Naturally Processed Peptides Bound To HLA-Dr Alleles" J. Exp. Med, 178, 27-47; (1993)														
1			Sctte, A. et al., "Capacity Of Intact Proteins To Bind To MHC Class II Molecules" J. Immunol., 143, 1265-1267; (1989)											
			1	Brown, J.H., et al. "Three-Dimensional Structure Of The Human Class II Histocompatability Antigen HLA-DRI Nature, 364, 33-39 (1993)										
				Kwon, et al., "Isolation And Sequence Of A cDNA Clone For Human Tyrinsinanse That Maps At The Mouse c-Albino Locus" PNAS, 84:7473-7477, (1987)										
			Shih	ihara,	S. et a	i., "M	ofecul	ar Bas	is For The Heter	ogeneity Of Human Ty	rosinase" J.	Exp. Med, 156:40	3-414, (1988)	
				Sette, A. et al., "HLA-DR4w4-Binding Morifs Illustrate The Bio chemical Basis Of Degeneracy And Specificity In Peptide-Dr Interactions" J. Immunol., 151:3163-3170, (1993)										
			Rair	Rammenscc, H.G. et al., "MIIC Ligands And Peptide Motifs: First Listing" Immunogenetics, 41:178-228 (1995)										
				Kozono, H. et al., "Production Of Soluble MHC II Class II Proteins With Covalently Bound Single Peptides" Nature, 369:151-154 (1994)										
			Sini	Sinigaglia, F. et al., "Motifs And Super motifs for MHC Class II Binding Peptides" J. Exp. Med., 181:449-451 (1995)										
V	Kawakami, et al. "Cloning Of The Gene Coding For A Shared Human Melanoma Antigen Recognized By Autologous T-Cells Infiltrating Into Tumor" Proc. Natl Acad. Sci., 91:3575-3579 (1994)													

Shees 2 of 4

FORM PTO-1449	U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTY, DECEMBER NO. 2026–4205US	SERIAL NO. FAX RECEI 08/533,895						
INFORMATIO	ON DISCLOSURE CITATION		NOV 0 3 19	98					
(Use several sheets if necessary)		Ti.	GROUP 16	300					
(036 36	verui sneets ij necessary)	APPLICANT(S) Topalian, et al.							
		FILING DATE September 26, 1995	GROUP ART LINIT 1645						
Wang, R-F et al., "Identification Of Infiltrating Lumpitocytes" J. Exp. M.		f A Gene Encoding A Melanoma Tumor Antigen Recognized By HLA-A31-Restricted Tumor Med., 181:799-804, (1995)							
		Sidney, J. et al., "DR B1* 0301 Molecules Recognize A Strentural Motif Distinct From The One Recognized By Most DR B, Alleles" J. Immunol., 149, 2634-2640, (1992)							
	Malcherek, G. et al., "Super Motifs Histocompatability Complex-class II		rived Peptides To Interact With Many Major 17-536 (1995)						
	Topalian, S., "MHC Class II Restrict in Immunology, 6:741-745, (1994)	tricted Tumor Antigens And The Role Of CD4+ T Cells In Cancer Antigens* Current Opinion							
		tian, S., et al. (1994) "Melanoma Specific CD4 + T-Lymphocytes Recognize Human Melanoma Antigens Processed And need By Epstein Barr Virus Transformed Cells" Int. J. Cancer 58:69-79							
	Nanda, et al., "Induction of Anti Sel	nda, et al., "Induction of Anti Self-Immunity to Bone Cancer" Cell, 82:13-17, (1995)							
		Markus, N. ct al., "Analysis of Cylokine Secretion by Melanoma-Specific CDY & Lymphocytes" Journal of Interferon and Cylokine Research, 15:739-746, (1995)							
	GenBank, Acession No. 103581 - Ja	enBank, Acession No. 103581 - January 14, 1995							
	GenBank, Accession No. U01873 - Se	873 - September 27, 1993							
	GenBank, Accession No. Y00819 - In	GenBank, Accssion No. Y(0)819 - July 28, 1995							
	GenBank, Accssion No. M27160 - I	GenBank, Accssion No. M27160 - December 2, 1996							
	Coulie, P.G. et al. (1993) "Genes co 14:104-109.	Conlie, P.G. et al. (1993) "Genes coding for tumor antigens recognized by human cytolytic T-lymphocytes." <i>J. Immunotherap.</i> ; 14:104-109.							
		Coulie P.G. et al. "A new gene coding for a differentiation antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas." J. Exp. Med. 1994; 180:35-42.							
,	Marcsh, C.A. et al.: "Cloning and o 1994; 13:87-95.	Marcsh, C.A. et al.: "Clouing and expression of the gene for the melanoma associated ME20 antigen." DNA and Cell Biology 1994; 13:87-95.							
	Cox, A.L., et al. "Identification of a 264:716-719.	peptide recognized by five melan	oma-specific human cytotoxic T cell lines." Science	ce 1994;					
	Brichard, V., et al.: "The tyrosinass melanomas". J. Exp. Med. 1993; 17		ized by autologous cytolytic T lymphocytes on Hl	LA-A2					
	Gaugier, B., et al. "Human gene Ma lymphocytes". J. Exp. Med. 1994;	AGE-3 codes for an antigen recogn 179:921-930.	ized on a melanoma by autologous cytulytic T						
	Traversari, C., et al.: "A nonapopti directed against tumor antigen M/2-		-1 is recognized on HLA-A1 by cytolytic T lymp 1457.	hocytes					
	Cellis, E., et al.: "Induction of anti- peptides epitopes". Proc. Natl. Acad	Cellis, E., et al.: "Induction of anti-turnor cytotoxic T lymphocytes in normal humans using primary cultures and synthetic peptides epitopes". <i>Proc. Natl. Acad. Sci. USA</i> 1994; 91:2105-2109.							
	Boon, T.: "Toward a genetic analys	on, T.: "Toward a genetic analysis of tumor rejection antigens". Adv. Cancer Res. 1992; 58:177-210.							
	Kawakami, Y., ct al.: "I'-cell recog	Kawakami, Y., ct al.: "I'-cell recognition of human melanoma antigens." J. Immunother. 1993; 14:88-93.							
		Bakker, A.B.H., et al.; "Melanocyte lineage-specific antigen gp100 is recognized by melanocyte-derived tumor infiltrating lymphocytes." J. Exp. Med. 1994; 179:1005-1009.							
	Wölfel, T., et al.: "Two tyrosinasc Eur. J. Immunol. 1994; 24:759-764.	nonapeptides recognized on HLA-	A2 melanomas by autologous cytolytic T. lympho	cytes."					

Short 3 of 4

		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTY, DOCKET NO. 2026-4205US	SERIAL NO. 08/533,895				
INFORMATION DISCLOSURE CITATION								
(Use several sheets if necessary)								
			APPLICANI(S) Topalian, et al.					
			FILING DATE September 26, 1995	GROUP ART UNIT 1645				
QKM		Adema, G.J., et al.: "Melanucyte lin HMB-45 are encoded by a single cibi	ineage-specific antigens recognized by monoclonal antibatics NK1-betch, HMB-50, and NA." Am J. Pathol. 1993; 143:1579-1585.					
		Kwon, B.S., et al.: "A melanocyte-s is in a syntenic region on human chro	Kwon, B.S., et al.: "A melanocyte-specific gene, Pmel 17, maps near the silver coat color locus on mouse chromosome 10 ar is in a syntenic region on human chromosome 12." <i>Proc. Natl. Acad. Sci. USA</i> 1991; 88:9228-9232.					
		Rosenberg, S.A., et al.: "Use of tun metastatic melanoma. Preliminary re	Rosenberg, S.A., et al.: "Use of tumor infiltrating lymphocytes and interleukin-2 in the immunotherapy of patients with metastatic melanoma. Preliminary report." N. Engl. J. Med. 1988; 319:1676-1680.					
		Kawakami, Y., ct al.,: "Shared hum transfected melanomas." J. Immunoi	nan melanuma antigens. Recognition by I 1992; 148:638-643.	y tumor infiltrating lymphocytes in HLA-A2.1				
		Van der Bruggen, et al.: "A gene en 1991; 254:1643-1647.	Van der Bruggen, et al.: "A gene encoding an antigen recognized by cytolytic T. lymphocytes on a human melanoma." Sci 1991; 254:1643-1647.					
		Falk, K., et al.: "Allele-specific mot 351:290-296.	Falk, K., et al.: "Allele-specific motifs revealed by sequencing of self-peptides eluted from MHC molecules." Nature 1991; 351:290-296.					
		Kubo, R., et al.: "Definition of spec 3924.	Kubo, R., et al.: "Definition of specific peptide motifs for four major HLA-A Alleles." Journal of Immunology 1994, 152:3913-3924.					
		Parker, K., et al.: "Sequence motifs 1992; 3580-3587.	Parker, K., et al.: "Sequence motifs important for peptide binding to the human MHC class 1 molecule. HLA-A2." J. Immunol. 1992; 3580-3587.					
		Ruppert, J., et al.: "Prominent role o 74:929-937.	Ruppert, J., et al.: "Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules." Cell 1993; 74:929-937.					
		Storkus, W., et al.: "Identification of lymphocytes." Journal of Immunolog	Storkus, W., et al.: "Identification of human melanoma peptides recognized by class 1 restricted tumor infiltrating T lymphocytes." Journal of Immunology 1993; 151:3719-3727.					
		Kawakami, Y., et al.: "Cloning of the infiltrating into tumor." Pro. Natl. A	Kawakami, Y., et al.: "Cloning of the gene coding for a shared human melanoma antigen recognized by autologous T cells infiltrating into tumor." Pro. Natl. Acad. Sci. USA 1994; 91:3515-3519.					
		Adema, G.J. et al., "Molecular chara Chemistry 1994; 269:20126-20133.	Adema, G.J. et al., "Molecular characterization of the melanocyte lineage-specific antigen gp100." Journal of Biological Chemistry 1994; 269:20126-20133.					
		EMBL DATABASE ACESSION NU MRNA 3'end' - June 15, 1990	EMBL DATABASE ACESSION NUMBER M32295; 26-11-90 Vogel A.: Human KD melanocyte specific secreted glycoprotein MRNA 3'end' - June 15, 1990					
		Kawakami, Y., et al.: "Identification of a human melanoma amigen recognized by tumor-infiltrating lymphocytes associated with In vivo tumor rejection." Proc. Natl. Acad. Sci. USA 1994; 91:6458-6462.						
		Kawakami, Y., et al., "Identification the Majority of HLA-A2-restricted To	Kawakami, Y., et al., "Identification of the Immunodominant Peptides of the MART-1 Human Melanoma Autigen Recognized by the Majority of HLA-A2-restricted Tumor Infiltrating Lymphocytes" J. Exp. Med. 180:347-352, 1994					
		Melanoma Patients by In Vitro Stimul	Rivoltini, L., et al., "Induction of Tumor-Reactive CTL from Peripheral Blood and Tumor-Infiltrating Lymphocytes of Melanoma Patients by In Vitro Stimulation with an Immunodominant Peptide of the Human Melanoma Antigen MART-1"  Journal of Immunology, 1995, 154:2257-2265					
		Slingluff, C.l., Jr., et al., "Direct and 740	Slingluff, C.1., Jr., et al., "Direct analysis of tumor-associated peptide antigens" Current Opinion in Insumulology 1994, 6:733-740					
		Cole, D.J., et al., "Characterization of MART-I Melanoma Autigun" Cancer	Cole, D.J., et al., "Characterization of the Functional Specificity of a Cloned T-Cell Receptor Heterodimer Recognizing the MART-I Melanoma Antigen" Cancer Res. 55:748-752 Feb. 1995					
		Cole, D.J., et al., "Identification of N Joining Regions Recognize the Same"	Cole, D.J., et al., "Identification of MART-1-specific T-Cell Receptors: T Cells Utilizing Distinct T-Cell Receptor Variable and Joining Regions Recognize the Same Tumor Epitope" Cancer Res. 54:5265-5268, 1994					
A		Castelli, C., et al., "Mass Spectrometric Identification of a Naturally Processed Melanoma Peptide Recognized by CD8."  Cytotoxic T Lymphocytes" J. Exp. Med. 181:363-368 1995						

Sheet 4 of 4

FORM PTO-1449 U.		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTY, DUCKET NO. 2026-4205US	SERIAL NO. 08/533,895					
INFOR	RMATION	DISCLOSURE CITATION							
	(Use sever	ral sheets if necessary)							
			APPLICANI(5) Topalian, et al.						
<del>.</del>			FILING DATE September 26, 1995	GROUP ART UNIT 1645					
TKN		Scnc, A., ct al., "Peptide Unding To Assays" Molecular Immunology 31:8	o The Most Frequent HLA-A Class I Alleles Measured By Quantitative Molecular Binding 813-822, 1994						
		Wölfel, T., et al., "Analysis Of Anti (CTL)" Int. J. Cuncer 55:237-244, 1	gens Recognized On Human Molanonu 993.	elanoma Cells By A2-Restricted Cytolytic T Lymphocytes					
		Wölfel, T., et al., "Isolation Of Natu Melanoma Cells In Association With	Wölfel, T., et al., "Isolation Of Naturally Processed Peptides Recognized By Cytolytic Lymphocytes (CTL) On Human Melanoma Cells In Association With HLA-A2.1" Int. J. Cancer 57:413-418, 1994.						
		Topalian, S.1., et al., "Human CD4" Tyrosinase Gene" PNAS 91:9461-946	Topalian, S.I., et al., "Human CD4 <sup>+</sup> T Cells Specifically Recognize a Shared Melanoma-Associated Antigen Encoded by the Tyrosinase Gene" <i>PNAS</i> 91:9461-9465, 1994.						
		Boël, P., et al., "BAGE: a New Gene Immunology 2:167-175 1995	Buel, P., et al., "BAGE: a New Gene Encoding an Antigen Recognized on Human Melanomas by Cytolytic T Lymphocytes" Immunology 2:167-175 1995						
		Slingluff, C.L., Jr., et al., "Recognit Mediated by at Least Six Shared Pept	Slingluff, C.L., Jr., et al., "Recognition of Human Melanoma Cells by HLA-A2.1-Restricted Cytotoxic T Lymphocytes Is Mediated by at Least Six Shared Peptide Epitopes" Journal of Immunology 150:2955-2963 1993						
		GENBANK DATABASE ACCESSION	GENBANK DATABASE ACCESSION NUMBER M77348 - Human PMEL 17 in RNA - Neventher 14, 1991 January 8, 195						
		GENBANK DATABASE ACCESSIO	GENBANK DATABASE ACCESSION NUMBER 1706654 - Human Differentiation Antigen Melan-A Protein in RNA - July 30.						
		GENBANK DATABASE ACCESSIO MRNA - June 25, 1994	GENBANK DATABASE ACCESSION NUMBER 1/06452 - Human Melanoma Antigen Recognized by T-Cells (MART-1) MRNA - June 25, 1994						
		GENBANK DATABASE ACCESSIO January 25, 1995	GENBANK DATABASE ACCESSION NUMBER \$73003 - GP100 Melanocyte Lineage Specific Antigen / PMELL 7 January 25, 1995						
		GENBANK DATABASE ACCESSION	ON NUMBER U01874 - Human ME20	:20 MRNA May 27, 1994					
		Bouchard, Brigitte, et al.; "Induction 1989; 169:2029-2042.	Bouchard, Brigitte, et al.: "Induction of pigmentation in mouse fibroblasts by expression of human tryosinance." J. Exp. Med. 1989; 169:2029-2042.						
		Tripathi, Ram K., et al.: "Tyrosinase Clusts of Missense Substitutions." An	Tripathi, Ram K., et al.: "Tyrosinase Gene Mutations in Type I (Tyrosinase-Deficient) Oculocutaneous Albinism Define Two Clusts of Missense Substitutions." American Journal of Medical Cenetics 1992; 43:865-871.						
		Octting, William S. and King, Richar Mutations and Polymorphisms of the	Oetting, William S. and King, Richard A.: "Molecular Basis of Type I (Tyrosinase-Related) Oculocutaneous Albinism: Mutations and Polymorphisms of the Human Tyrosinase Gene." Human Mutation 1993; 2:1-6.						
		Spritz, Richard A.: "Molecular Gene	Spritz, Richard A.: "Molecular Genetics of Oculocutaneous Albinism." Seminars in Dermatology 1993; Vol. 12, No.3:167-172						
		Robbins, et al. (1994), "Recognition of immunotherapy", Cancer Research; 5	Robbins, et al. (1994). "Recognition of tyrosinase by tumor-infiltating lymphocates from a patient responding to immunotherapy", Cancer Research; 54:3124-3126						
		Robbins, et al (1995), "Cloning of a new gene encoding an antigen recognized by melanoma-specific HLA-A24-restricted turnor-infiltrating lymphocytes" <i>J. IMMUNOL.</i> 1995;154(11), 5944, 50							
EXAMINER	5		DATE CONSIDERED	DATE CONSIDERED 1-18-99					
EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not									